

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

SNYDER *et al.*

Appl. No.: 09/849,781

Filed: May 4, 2001

For: **PROTEIN CHIPS FOR HIGH-
THROUGHPUT SCREENING OF
PROTEIN ACTIVITY**

Confirmation No.: 9891

Art Unit: 1639

Examiner: Wessendorf, Teresa D.

Atty. Docket: 2493.0010002/RWE/JKM

Declaration of Michael Snyder Under 37 C.F.R. § 1.132

Mail Stop Amendment

Commissioner for Patents

PO Box 1450

Alexandria, VA 22313-1450

Sir:

The undersigned, Michael Snyder, residing at 230 Somerset Avenue, Fairfield, CT 06824, USA, declares and states as follows:

1. I am a co-inventor of above-captioned U.S. Patent Application No. 09/849,781 ("the '781 application") entitled "Protein Chips for High-Throughput Screening of Protein Activity," which has a filing date of May 4, 2001, which claims benefit to U.S. Provisional Patent Application No. 60/201,921, filed May 4, 2000, and to U.S. Provisional Patent Application No. 60/221,034, filed July 27, 2000.

2. I am currently employed by Yale University, the assignee of the above-captioned application. I hold the positions of Director of the Yale Center for Genomics and Proteomics and Professor in the Departments of Molecular, Cellular and

Atty. Dkt. No. 2493.0010002/RWE/JKM

Developmental Biology, and Molecular Biophysics and Biochemistry. My credentials are provided in the *curriculum vitae* that is attached to this declaration as Exhibit A. I received my Ph.D. degree in Biology from the California Institute of Technology in 1983, where my research was focused on *Drosophila* gene organization. As seen from my attached *curriculum vitae*, I have published many papers related to characterization of yeast and human proteomes, control of cell division and cellular morphology in yeast, and characterization of the human genome. I have received many honors and am involved in numerous professional and scientific societies related to genomics, proteomics and genetics. Based on my education and experience, I am an expert in the field of yeast and human genomics, proteomics, and molecular genetics.

3. I have reviewed and am familiar with the '781 application filed on May 4, 2001, the Office Action dated March 18, 2008 ("the Office Action"), issued by the U.S. Patent and Trademark Office in the present application, the references cited in the Office Action, and the currently pending claims, filed in the Reply to Office Action with this declaration.

4. In the Office Action, the Examiner asserts that the claims are allegedly not enabled. Specifically, the Examiner asserts that the specification does not enable the claimed array comprising 61 purified, active kinases and functional domains of kinases, from a mammal, yeast or *Drosophila*.

5. In making this declaration, it is my opinion that at the time this application was filed, a typical practitioner in the field of genomics and proteomics would have been able to make and use the claimed positionally addressable arrays based

on knowledge available to those in the field in combination with the disclosure of the '781 application. It is also my opinion that any experimentation required for making and using the claimed positionally addressable arrays would have been routine and thus not inordinate or excessive. Therefore, it is my opinion that the rejection is in error, and I present herein the rationale upon which I base that conclusion.

6. As discussed in detail below, the specification of the '781 application clearly provides sufficient disclosure for a typical practitioner in the field of proteomics to make and use positionally addressable arrays comprising 61 purified, active yeast kinases. Based upon this disclosure, in combination with what was known at the time of filing of the present application, the use of kinases from other organisms, including mammals and *Drosophila*, in the preparation of the presently claimed positionally addressable arrays, would not have required undue experimentation, but rather, routine and straightforward experiments.

7. As set forth in the references discussed below, protein kinases and functional kinase domains used in the positionally addressable arrays that form the basis of the claims were, at the time this application was filed, all well-known, well-characterized proteins. For example as discussed in Hunter and Plowman, "The protein kinases of budding yeast: six score and more," *TIBS* 22:18-22 (1997) (hereinafter "Hunter and Plowman"):

[b]udding yeast has 113 convention protein kinase genes, corresponding to ~2% of the total genes (see Table 1 in cetrefold). More than 60% of these protein kinases have either known or suspected functions; the remainder are novel, and functional analysis awaits. In terms of defined functions encoded by the yeast

genome, protein kinases come in a close second behind transcription factors.

Hunter and Plowman at page 18, first column, first paragraph. A review by Manning *et al.*, "The Protein Kinase Complement of the Human Genome," *Science* 298:1912-1934 (2002) (hereinafter "Manning1"), indicates "[p]rotein kinases are among the largest families of genes in eukaryotes and have been intensively studied." Manning1 at page 1913, first column, first paragraph.

8. It is well recognized that kinases are highly conserved such that homologs exist between yeast and many other organisms. Furthermore, the regulation of the different kinases and the phosphorylation motifs of substrates recognized by related kinases are often the same, indicating that they behave similarly biochemically. As set forth in Manning *et al.*, "Evolution of protein kinase signaling from yeast to man," *TRENDS in Biochemical Sciences* 27:514-520 (2002) (hereinafter "Manning2"):

all major kinase groups and most kinase families are shared among metazoans, and many are also found in yeast, reflecting the breadth of conserved function mediated by kinases. This ancient conservation enables cross-species analysis of function, particularly of human kinases in simpler model systems. Of 209 subfamilies, 51 are present in all four genomes, and 144 are present in all metazoans, indicating that most divergence of kinases into specific functions and families occurred during early eukaryotic and metazoan evolution.

Manning2 at page 514, second column, first full paragraph, and at Figure 1 and Table 1.

This known relationship is also addressed in Manning 1:

Phylogenetic comparison of the human kinome with that of yeast, worm, and fly confirms that most kinase families are shared among metazoans and defines classes that are expanded in each lineage. Of 189 subfamilies present in human, 51 are found in all four eukaryotic kinomes, and these

presumably serve functions essential for the existence of a eukaryotic cell. An additional 93 subfamilies are present in human, fly, and worm, implying that these evolved to fulfill distinct functions in early metazoan evolution. Comparison with the draft mouse genome indicates that more than 95% of human kinases have direct orthologs in mouse; additional orthologs may emerge as that genome sequence is completed.

Manning¹ at page 1914, last paragraph bridging first and second columns. *See also* Hunter and Plowman at page 20, second column, second paragraph, "[m]ost of the main vertebrate subfamilies of protein kinases are represented in yeast."

9. Furthermore, as function is often highly conserved, human kinases can be substituted for yeast kinases, illustrating the highly conservative nature of these proteins. *See* Lee and Nurse, "Complementation used to clone a human homologue of the fission yeast cell cycle control gene *cdc2*," *Nature* 327: 31-35 (1987) (hereinafter "Lee and Nurse"):

Because the human *CDC2Hs* gene can provide all of the functions of *cdc2Sp* in fission yeast it is reasonable to assume that it performs a similar role to *cdc2Sp* in controlling the human cell cycle. This conclusion is supported by the structural similarity between the two genes both in overall homology and size of the proteins. . . .

The identification of a *cdc2*-like function in human cells suggests that elements of the mechanisms by which the cell cycle is controlled will probably be found in all eukaryotic cells.

Lee and Nurse at page 35, first column to second column.

10. Therefore, at the time this application was filed, the "state of the art" in protein kinases was such that a practitioner possessing a typical level of skill in proteomics, such skill including but not limited to protein purification and analysis,

would have readily recognized from the '781 application, and the knowledge available in the art, that kinases of yeast, mammals and *Drosophila* could be utilized to practice the present invention.

11. The Examples set forth in the '781 application describe positionally addressable protein arrays made with purified, active kinases isolated from yeast. The specification describes that the purified, active yeast kinases were prepared by cloning yeast kinase genes into a high copy *URA3* expression vector. See the '781 application at page 26, lines 27-28. The plasmids containing the vector sequences were transformed into yeast, and Ura⁺ colonies were selected. Plasmids were rescued in *E. coli*, then transformed into the *pep4* yeast strain for kinase protein purification. Purified, active kinases were attached to polydimethylsiloxane (PDMS) chips, and the chips comprising the purified, active yeast kinases were assayed for the phosphorylation of 17 different substrates to determine *in vitro* kinase activity. This large-scale analysis of purified, active yeast kinases allowed me to characterize the activities of 119 protein kinases for 17 different substrates and to compare the functional relationship of these protein kinases to one another. This analysis also allowed me to determine meaningful biological interactions between these kinases and their targets.

12. As described above, the '781 application provides working examples where purified, active kinases from yeast were used to make a positionally addressable array. As described in the references cited above, those working in the field of proteomics at the time this application was filed were aware that yeast, mammalian and *Drosophila* kinases are highly conserved and homologous. See e.g., Manning2. In

addition, the working examples in the specification provide a clear road map sufficient to allow a practitioner in the field of genomics and proteomics to extend the experiments with yeast kinases to mammalian and *Drosophila* kinases making such an extension routine. *See, e.g.*, the '781 application at pages 25-35. In particular, in my opinion, a typical practitioner in the field of proteomics would consider the production of an array using at least 61 purified, functional kinases from yeast as detailed in the specification sufficient to enable them to prepare arrays using purified, active kinases from other organisms, including mammals and *Drosophila*, especially in light of the specific reference to this use in the specification.

13. The Examiner has also rejected claims 1-11, 141, 181-186, 188 and 193-195 as allegedly being obvious in view of Uetz *et al.*, *Nature* 403:623-631 (February 10, 2000) (hereinafter "Uetz"). The Examiner contends that Uetz discloses a protein array comprising yeast genome encoded proteins, and that the proteins were expressed in 96-well plates. The Examiner asserts that the claimed kinases would have been inherent to the yeast array disclosed in Uetz, since yeast inherently contain kinase in their structure, or that they would have been obvious to determine given the identified genome of yeast. It is my opinion that there is no disclosure in Uetz, inherent or otherwise, of an array or the construction of an array of at least 61 kinases or functional kinase domains, in which the array comprises kinases that are *purified and active*, as required by present claim 1. At the time of filing of the present application, as set forth below, it was unexpected that kinases and functional kinase domains of these kinases, could be purified and placed on a solid support to form an array, and that these kinases and kinase domains, *would retain their kinase activity*. Skeptics in the field, before, and even well after the time the

present application was filed, thought that the proteins used to make the array of the presently claimed invention would denature and, therefore, be inactive. It was an unexpected and surprising result that the purified proteins on the array retained their activity and could be utilized to determine meaningful biological interactions between the purified, active kinases and their targets (such as enzyme-enzyme or enzyme-substrate interactions). Simply put, it is only after the guidance provided in the present specification that a practitioner in the field of proteomics would consider it possible to generate the presently claimed arrays.

14. The following exemplary references describe the skepticism from those in the field regarding the preparation of protein arrays both before and after the time of filing of the present application, as well as some of the problems regarding preparation of protein arrays comprising large numbers of purified active proteins that were overcome by the presently claimed invention.

15. Anderson, K.S. and LaBaer, J., *Journal of Proteome Research* 4:1123-1133 (March 30, 2005) (hereinafter "Anderson and LaBaer") state that

[t]heir theoretical advantages notwithstanding, protein microarrays have still not found widespread use, in part because producing them is challenging. Historically, it has required the high-throughput production and purification of protein, which then must be spotted on the arrays. Once printed, concerns remain about the shelf life of proteins on the arrays.

Anderson and LaBaer, page 1129.

16. Shaw, G., *Drug Discovery and Development* (February 3, 2005) (hereinafter "Shaw") states that:

[i]t was first thought that protein biochips would just be an extension of DNA microarrays, and that hasn't exactly panned out," says Bodovitz. That's because proteins have proven to be much trickier to work with in array format than their genomic counterparts. First of all, there are issues of stability. Membrane proteins, for example, make up the majority of potential drug targets, but they're particularly challenging to stabilize. Then there's the choice of immobilization technique, which determines how well the target protein presents itself to the capture agent, and the problem of nonspecific binding. And of course, proteins are inherently unstable outside their natural habitat of living cells, making them much more challenging than DNA to tag and manipulate.

Shaw at page 1.

17. As set forth in the following references, it was well known that spotting proteins on solid surfaces often resulted in protein denaturation caused by uncontrolled absorption, thereby inactivating the proteins. See Abstract of both Tleugabulva *et al.*, "Evidence for the denaturation of recombinant hepatitis B surface antigen on aluminum hydroxide gel," *J. Chromatogr. B. Biomed. Sci.* 702:153-163 (1998); and Servagent-Noinville *et al.*, "Conformational Changes of Bovine Serum Albumin Induced by Absorption on Different Clay Surfaces: FTIR Analysis," *J. Colloid Interface Sci.* 221:273-283 (2000).

18. Prior to the presently claimed invention, only small numbers of proteins, and/or inactive proteins, were able to be displayed on an array. It is only after the teaching of the '781 application that large numbers of purified proteins were able to be placed on a solid support, in the recited density, such that they remained active. In Bussow *et al.*, "A method for global protein expression and antibody screening on high-density filters of an arrayed cDNA library," *Nucleic Acids Res.* 26: 5007-5008 (1998) (hereinafter "Bussow1"), large scale protein arrays were produced, but only using

denatured proteins. ("These protein filters were processed on pre-soaked blotting paper, i.e., denatured in 0.5 M NaOH, 1.5 M NaCl for 10 min, neutralized for 2x5 min in 1 M Tris-HCl, pH 7.5, 1.5M NaCl and incubated for 15 min in 2x SSC. Filters were air-dried and stored at room temperature." Bussow1 at page 507, second column, first paragraph.) Similarly, in Bussow *et al.*, "A human cDNA library for high-throughput protein expression screening," *Genomics* 65:1-8 (2000) (hereinafter "Bussow2"), denaturing conditions were used to prepare the protein arrays. ("Twenty-five microliters of 50% Ni-NTA agarose was added to protein extracts obtained under denaturing conditions." Bussow2 at page 2, first column, third paragraph.) In addition, in the introduction of Bussow2, the authors discuss the difficulties in producing large amounts of purified proteins and maintaining their activity when placed on an array:

the individuality of protein molecules demands highly customized procedures for their expression. Automation of these procedures requires systems that allow the efficient handling of large numbers of clones representing many different proteins. Bacterial systems are easy to manage *but the expression of eukaryotic proteins can be problematic*, due to aggregation, formation of insoluble inclusion bodies, and/or degradation of the expression product. Eukaryotic systems suffer from lower yields of heterologous protein (e.g. *Saccharomyces cerevisiae*s), high demands on sterility (e.g. mammalian systems) or time consuming cloning procedures (e.g. Baculovirus system).

Bussow at page 1, last paragraph bridging first and second columns (citations emitted, emphasis added). Thus, rather than attempt to overcome these various issues, the authors proceeded to produce denatured proteins in *E. coli* for use in their arrays. Similarly, Lueking *et al.*, "Protein microarrays for gene expression and antibody screening," *Anal. Biochem.* 270:103-111 (1999) (hereinafter "Lueking"), also prepared protein arrays

utilizing denaturing conditions (6M guanidinium-HCl, 0.1M NaH₂PO₄, 0.01 M Tris-HCl, pH 8.0, Lueking at page 104, second column, first paragraph).

19. In Ge, "UPA, a universal protein array system for quantitative detection of protein-protein, protein-DNA, protein-RNA and protein-ligand interactions," *Nucleic Acids Research* 28:e3(i-vii), the author was only able to produce arrays comprising 48 proteins at a very low density, utilizing a traditional purification format. Extension of this disclosure to arrays comprising at least 100 different substances per cm² would have required extensive, additional experimentation beyond the scope of the disclosure provided in this reference.

20. Thus, as noted in the references cited above, it is my opinion that experts in the field were clearly skeptical of the ability to prepare protein arrays comprising large numbers of purified, active enzymes, at the densities recited in the presently claimed invention. Instead, denaturing conditions or traditional purification methods on small numbers of proteins were required. Absent the teachings of the '781 application there was no reasonable expectation that arrays comprising at least 100 different substances per cm², wherein the substances comprise 61 purified, active kinases on a solid support, could be successfully prepared, at the time of filing of the present application.

21. It is only after the guidance of the present specification, specifically the methods described in the present application which allowed for large-scale purification and arraying of active kinases or functional kinase domains, that preparation of the presently claimed positionally addressable arrays was possible. The methods of the present invention, as discussed below, are suitable for rapidly purifying large numbers of

samples, such as the purification of active kinases isolated from mammalian, yeast, and *Drosophila*. As an example, yeast kinases were purified by growing different strains of *pep4* yeast cells, each containing a plasmid encoding a single GST-tagged kinase gene, in 96-well plates. Galactose was added to induce protein expression. The cultures of the same strain were combined, washed, resuspended and lysed. The GST fusion proteins were purified from these strains using glutathione beads and standard protocols in a 96-well format. This method allowed for the purification of a high number of yeast kinases in a relatively short amount of time. The buffers and methods used also ensured that the purified kinases or functional kinase domains retained their activity. See Example 1 of the '781 application at page 26, line 25, through page 27, line 19.

22. Positionally addressable arrays were then prepared using the purified, active proteins purified as described above. Arrays were made from polydimethylsiloxane (PDMS) (Dow Chemical, USA), which was cast over microfabricated molds. Liquid PDMS was poured over the molds and, after curing flexible silicone elastomer array sheets were then peeled from the reusable molds. Arrays were immersed in 3-glycidooxypropyltrimethoxysilane linker (GPTS) in order to facilitate adsorption of protein to the wells. To attach proteins to the chips, protein solutions were added to the wells and incubated on ice for 1 to 2 hours. After rinsing with cold HEPES buffer, the wells were blocked with BSA in PBS. See Example 1 of the '781 application as filed at page 27, line 21, through page 28, line 9.

23. The methods set forth in the present application are sufficient to enable a typical practitioner in the field of proteomics to make and use positionally addressable

arrays comprising 61 purified, active kinases of yeast, mammals and *Drosophila*. Furthermore, such methods were only available upon filing of the present application.

24. The Examiner has also rejected claims 1-11, 141, 164, 169, 173, 174, 177, 181-186, 188 and 193-195 as allegedly being unpatentable over Shalon (WO 95/35505; hereinafter "Shalon") in view Felder *et al.* (U.S. Patent No. 6,458,533; hereinafter "Felder") of Lafferty (U.S. Patent No. 6,972,183; hereinafter "Lafferty").

25. The Examiner contends that Shalon discloses a microarray having regions with a density of at least about 100/cm², and that the arrays can comprise enzymes. The Examiner notes, however, that Shalon does not disclose arrays comprising kinases. The Examiner relies on the disclosures of Felder or Lafferty to cure this deficiency. Specifically, the Examiner contends that Felder discloses that kinases are enzymes, and the Lafferty discloses an array containing substrate-enzymes, such as kinases. The Examiner therefore concludes that it would have been obvious to prepare the array disclosed in Shalon using the kinases disclosed in Felder and Lafferty, and hence, the presently claimed invention is rendered obvious.

26. Shalon is primarily directed to arrays comprising polynucleotides (*see* Examples 1-3), and only mentions in passing that arrays comprising proteins and enzymes could be constructed. Furthermore, Felder discloses preparation of arrays comprising peptides that are *substrates* for kinases, not arrays comprising the kinases themselves, "[a] chimeric linker molecule is prepared in which a 25 base pair oligonucleotide complementary to one of the anchors is crosslinked to a *peptide substrate of a tyrosine phosphokinase enzyme*." Felder at column 44, lines 18-21

(emphasis added). Thus, Felder does not disclose the preparation of arrays comprising 61 purified active kinases or functional kinase domains thereof, as required by present claim 1.

27. With regard to Lafferty, it is important to note that the arrays disclosed therein are limited to enzymes expressed in expression library cells, and that Lafferty does not disclose the purification of these enzymes prior to placement on a solid support, as recited in the presently claimed invention. Lafferty states, at column 18, lines 1-14:

The library comprises a plurality of recombinant clones, which comprise host cells transformed with constructs comprising expression vectors into which have been incorporated nucleic acid sequences derived from the DNA samples. One or more substrates and at least a subset of the clones is then introduced, either individually or together as a mixture, into capillaries (all or a portion thereof) in a capillary array. Interaction (including reaction) of the substrate and a clone expressing an enzyme having the desired enzyme activity produces an optically detectable signal, which can be spatially detected to identify one or more capillaries containing at least one signal-producing clone. The signal-producing clones can then be recovered from the identified capillaries.

Therefore, Lafferty does not disclose arrays comprising 61 purified active kinases, as set forth in the presently claimed invention.

28. Therefore, it is my opinion that Shalon, Felder and Lafferty, alone or in combination, do not disclose the presently claimed positionally addressable arrays, specifically, arrays comprising 61 purified active kinases or functional kinase domains thereof, as set forth in present claim 1. Specifically, the references cited by the Examiner do not disclose preparation of purified kinases which are then placed on a solid support, as recited in present claim 1. Thus, there are clearly differences between the cited

references and the presently claimed invention that have not been addressed by the Examiner.

29. As set forth in detail above, there was no reasonable expectation of success of preparing an array comprising at least 61 purified active kinases or functional kinase domains of these kinases, based on the references cited by the Examiner. At the time of filing of the present application, experts in the field were skeptical, and it was unexpected, that purified kinases and functional kinase domains of these kinases, could be placed on a solid support to form an array, and that these proteins would retain their activity. It was also unexpected that these positionally addressable arrays could be used to determine meaningful biological interactions (such as enzyme-enzyme or enzyme-target interactions) between the purified active kinases and their targets. It is only after the guidance of the present specification that a typical practitioner in the field of proteomics would be able to generate the presently claimed arrays.

30. As discussed above, it is my opinion that, at the time of filing of the present application, it was unexpected that purified kinases and functional kinase domains of these kinases could be purified and placed on a solid support to form an array. It is also my opinion that it was unexpected that the purified kinases and functional kinase domains of these kinases would retain their activity when placed onto the array. It is only after the guidance provided in the present specification that a person of ordinary skill in the art would consider it possible to generate the presently claimed arrays.

31. I conclude that for at least the reasons stated above that there was no reasonable expectation of success of preparing an array comprising at least 61 purified active kinases or functional kinase domains of these kinases. It is my opinion that, at the time of filing of the present application, experts in the field were skeptical, and it was unexpected, that purified kinases and functional kinase domains of these kinases could be placed on a solid support to form an array, and that these proteins would retain their activity. It is only after the guidance of the present specification that a typical practitioner in the field of proteomics would be able to generate the presently claimed arrays.

32. I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the present patent application or any patent issued thereon.

Respectfully submitted,

Date: 4/13/09

A handwritten signature in black ink, appearing to read 'MSnyder', written over a horizontal line.

Michael Snyder

862691_1.DOC

CURRICULUM VITAE: Michael Snyder

Present Address Dept of Molecular, Cellular and Developmental Biology
 Yale University
 New Haven, CT 06520-8103
 Office (203) 432-6139
 Mobile (203) 376-2647
 michael.snyder@yale.edu

Education

1973-1977 B.A. University of Rochester, Rochester, New York
 Chemistry and Biology
 1978-1982 Ph.D. California Institute of Technology, Pasadena, California
 Department of Biology, Degree conferred 6/83

Professional Experience

1977-1978 Research Assistant, Department of Biology, University of
 Rochester, Rochester, New York
 1978-1982 Graduate Student with Dr. Norman Davidson, California Institute
 of Technology, Pasadena, California
 1982-1986 Postdoctoral Fellow with Dr. Ronald W. Davis, Department of
 Biochemistry, Stanford University School of Medicine,
 Stanford, California
 1986-1990 Assistant Professor, Department of Biology, Yale University,
 Connecticut
 1990-1997 Associate Professor, Department of Biology, Yale University,
 Connecticut (tenured 1994)
 1997-present Professor, Dept. of Biology (now Mol. Cell. Devel. Biol.), Yale Univ., CT
 1991-present Member Yale Comprehensive Cancer Center
 1992-present Associate Professor/Professor (Joint Appointment), Department of
 Molecular Biophysics and Biochemistry, Yale University
 1998-2004 Chair, Dept. of Molecular, Cellular and Developmental Biology, Yale Univ.
 (Dept. doubled in size and 3X in funds during my tenure).
 2002-present Director, Yale Center for Genomics and Proteomics
 2006-present Genetics Society of America Council Member (Elected)
 2006- 2008 President, US HUPO (Elected)

Academic Honors/Fellowships

1978-1982 NIH Predoctoral Training Fellowship
 1982-1985 Helen Hay Whitney Postdoctoral Fellowship
 1986 United Scleroderma Foundation Award
 1987-1991 Pew Scholar Award
 1989 Yale Junior Faculty Fellowship
 2000-2005 Burroughs Wellcome Scholar Award
 2002 Genome Technology Finalist in Microarray Masters
 2002-present Appointed Lewis B. Cullman Professor of MCDB
 2007 Connecticut Medal of Science

Advisory Committees

1989,90, 94-98, Member NIH Study Section- Ad hoc Reviewer

2000-07	
1993	Scientific Advisory Board Review Panel-American Cancer Society
1994-96	Howard Hughes Predoctoral Fellowship Review Committee
2002	Damon Runyan Walter Winchell Review Panel
1996-2002	March of Dimes Grant Review Panel
1997	ATCC Advisory Committee
1999	NSF Division Review Panel
2000, 2002	NIH Study Section Review Working Group
2000	Canadian Genome Center Review Panel
2001-present	Northeast Structural Genomics Consortium Scientific Advisory Committee
2001-present	Member, Institute of Genetics Advisory Council, CIHR Canada
2002-2007	Member, Ontario Genome Institute Scientific Advisory Board
2002-present	Member, Chinese National Human Genome Institute Advisory Board
2007-present	SAB, Integrated Genomics Project Univ. of Toronto
2008-present	SAB, Duke Univ. Systems Biology Center
2003-2006	SAB, Blueprint Initiative
2003	NIH Special Road Map Advisory Committee
2003	External Reviewer, Dept. of Medical Genetics, Univ. of Toronto
2003-present	Scientific Advisory Board, Gottenberg Univ. Genomics Meeting
2004-2007	Damon Runyan-Walter Winchell Review Panel
2007	NSF Plant Genomics Initiatives 5 Year Review
2008	External Review Committee, Gene Expression Unit, EMBL

Meeting Organizer

1994	Coorganizer, Juan March Meeting on Signal Transduction & Morphogenesis
1996	Chair, FASEB Meeting: Yeast Chromosome Structure, Repl. & Segregation
2000	Yeast Genetics Meeting Program Committee
2004	CoOrganizer, ASBMB Genomics, Proteomics and Bioinformatics Meeting*
2004	CoOrganizer, GSA International Meeting on Yeast Molecular Genetics
2004	CoOrganizer, CSH Plant Genomes: From Sequence to Phenomes
2006	CoOrganizer, ASBMB Genomics, Proteomics and Bioinformatics Meeting
2006	Organizer, GSA International Meeting on Yeast Molecular Genetics*
2006	CoOrganizer, US HUPO
2007	CoOrganizer, CSH Plant Genomes: From Sequence to Phenomes
2007	CoOrganizer, Keystone Meeting on Functional Genomics

Editorial Boards

1996-2000	Editorial Board, The Dynamic Cell
2001-2004	Editorial Board, Chemistry and Biology
2000-2006	Editorial Board, FEMS Yeast Research
2000-present	Editor-in-Chief (until 2002; now Editor) Functional and Integrative Genomics
2001-present	Editorial Board, Molecular and Cellular Proteomics
2002-present	Editorial Board, Drug Discovery Today
2004- present	Editorial Board, PloS Genetics
2005-present	Editorial Board, Genes and Development
2005-present	Editorial Board, Molecular Systems Biology

Named/Distinguished Lectureships (starting 2004)

2004	California Institute of Technology, Norman Davidson Lecture
2004	University of Chicago, Fredrick Setiz Lecture

2004 Tulane University, Gerber Lecture
 2004 University of Iowa, Raymond Fung Lecture
 2006 Northeastern University, Hoehn Lecture
 2007 EBI Distinguished Lecturer
 2008 Northwestern University Distinguished Lectureship
 2008 Fred Sherman Lecture, Univ. of Rochester

Keynote/Featured Speaker

2000 University of Sherbrooke Graduate Symposium*
 2001 CHI: Functional Genomics
 2002 Yale University Pathology Retreat
 2002 Georgia Bioinformatics Symposium
 2003 University of Texas Medical Faculty Symposium
 2003 Structural and Functional Genomics, Singapore
 2004 International Meeting On Arabidopsis*
 2004 Pennsylvania State Graduate Student Symposium
 2005 Keystone Conference on Plant Signaling
 2005 New York University: Genomics Symposium—Genomes in Action
 2005 Chip to Hits
 2005 Systems Biology Meeting
 2006 AGCT Genomics Meeting
 2006 Genomes to Biology Meeting, Manchester, UK
 2006 Keystone Conference on Plant Abiotic Stress
 2006 CHI Genes to Targets
 2006 AUHUPO*
 2007 ABRF*
 2007 Pan American and Brazilian Biochemistry and Molecular Biology Meeting*
 2007 Uppsala Neuroscience Center Launch
 2007 IBC Diagnostics 2 Discovery
 2007 DREAM 2 Conference
 2008 CHI Mining the Plasma Proteome
 2008 Student Invited Speaker, Vanderbilt Genetics Retreat
 2008 Genetics Day, Univ. of Rochester

*Meetings attended by over 900 people

Commercial Activity

Topogenetics/Exelexis 1990-1991 Partial Founder
 Genaissance 1999-2001 Scientific Advisory Board (SAB)
 Mycota 1996-2000 SAB
 Protometrix 2001-2004 (now subsidiary of Invitrogen) Founder and Chair of SAB
 RxGen 2003-present SAB
 Affomix 2006- Present Founder and Chair of SAB

Publications

1. Kung LA*, Tao S-C*, Qian J, **Snyder M**, Zhu H. Global analysis of the glycoproteome in *S. cerevisiae* reveals new roles for protein glycosylation in eukaryotes. Submitted.
2. Korb J, Kim PM, Chen X, Urban AE, Weissman S, **Snyder M**, Gerstein MB. The current excitement about copy-number variation: how it relates to gene duplications and protein families. *Curr Opin Struct Biol*. 2008. [Epub ahead of print] PMID: 18511261

3. Lian Z, Karpikov A, Lian J, Mahajan MC, Hartman S, Gerstein M, **Snyder M**, Weissman SM. A Genomics Analysis of RNA polymerase II modification and chromatin architecture related to 3' end RNA polyadenylation. *Genome Res.* 2008 May 16. [Epub ahead of print] PMID: 18487515.
4. Nagalakshmi U, Wang Z, Waern K, Shou C, Raha D, Gerstein M, **Snyder M**. The Transcriptional Landscape of the Yeast Genome Defined by RNA Sequencing. *Science.* 2008 May 1. [Epub ahead of print] PMID: 18451266
5. Wu JQ, **Snyder M**. RNA polymerase II stalling: loading at the start prepares genes for a sprint. *Genome Biol.* 2008 May 2;9:220. PMID: 18466645
6. Wu JQ, Du J, Rozowsky J, Zhang Z, Urban AE, Euskirchen G, Weissman S, Gerstein M, **Snyder M**. Systematic analysis of transcribed loci in ENCODE regions using RACE sequencing reveals extensive transcription in the human genome. *Genome Biol.* 2008;9: R3.
7. Nath AK, Brown RM, Michaud M, Sierra-Honigsmann MR, **Snyder M**, Madri JA. Leptin affects endocardial cushion formation by modulating EMT and migration via Akt signaling cascades. *J. Cell Biol.* 2008 181:367-80. Epub 2008 Apr 14. PMID: 18411306
8. Johnson DS, Li W, Gordon DB, Bhattacharjee A, Curry B, Ghosh J, Brizuela L, Carroll JS, Brown M, Flicek P, Koch CM, Dunham I, Bieda M, Xu X, Farnham PJ, Kapranov P, Nix DA, Gingeras TR, Zhang X, Holster H, Jiang N, Green R, Song JS, McCuine SA, Anton E, Nguyen L, Trinklein ND, Ye Z, Ching K, Hawkins D, Ren B, Scacheri PC, Rozowsky J, Karpikov A, Euskirchen G, Weissman S, Gerstein M, **Snyder M**, Yang A, Moqtaderi Z, Hirsch H, Shulha HP, Fu Y, Weng Z, Struhl K, Myers RM, Lieb JD, Liu XS. Systematic evaluation of variability in ChIP-chip experiments using predefined DNA targets. *Genome Res.* 2008, in press.
9. Chang W, Zaarour RF, Reck-Peterson S, Rinn J, Singer RH, **Snyder M**, Novick P, Mooseker MS. Myo2p, a class V myosin in budding yeast, associates with a large ribonucleic acid-protein complex that contains mRNAs and subunits of the RNA-processing body. *RNA.* 491-502.
10. Hudson ME, Pozdnyakova I, Haines K, Mor G, **Snyder M**. Identification of differentially expressed proteins in ovarian cancer using high-density protein microarrays. *Proc Natl Acad Sci USA.* 2007;104: 17494-9.
11. ENCODE Project Consortium. Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature.* 2007;447: 799-816.
12. Korbel JO,* Urban AE,* Affourtit J,* Godwin B, Grubert F, Simons JF, Kim PK, Palejev D, Carriero N, Du L, Taillon B, Tanzer A, Chi J, Yang F, Carter N, Hurles ME, Weissman S, Harkins T, Gerstein M, Egholm M, **Snyder M**. Paired-end mapping reveals extensive structural variation in the human genome. *Science.* 2007;318: 420-6.
13. Robertson G, Hirst M, Bainbridge M, Bilenky M, Zhao Y, Zeng T, Euskirchen G, Bernier B, Varhol R, Delaney A, Thiessen N, Griffith OL, He A, Marra M, **Snyder M**, Jones S. Genome-wide profiles of STAT1 DNA association using chromatin immunoprecipitation and massively parallel sequencing. *Nat Methods.* 2007;4: 651-7.
14. Zhu X, Gerstein M, **Snyder M**. Getting connected: analysis and principles of biological networks. *Genes Dev.* 2007;21: 1010-24.

15. Zhang ZD, Rozowsky J, Lam HY, Du J, **Snyder M**, Gerstein M. Telescope: online analysis pipeline for high-density tiling microarray data. *Genome Biol.* 2007;8: R81.
16. Korbel JO, Urban AE, Grubert F, Du J, Royce TE, Starr P, Zhong G, Emanuel BS, Weissman SM, **Snyder M**, Gerstein MB. Systematic prediction and validation of breakpoints associated with copy-number variants in the human genome. *Proc Natl Acad Sci USA.* 2007;104: 10110-5.
17. Gerstein MB, Bruce C, Rozowsky JS, Zheng D, Du J, Korbel JO, Emanuelsson O, Zhang ZD, Weissman S, **Snyder M**. What is a gene, post-ENCODE? History and updated definition. *Genome Res.* 2007;17: 669-81.
18. Trinklein ND, Karaoz U, Wu J, Halees A, Force Aldred S, Collins PJ, Zheng D, Zhang ZD, Gerstein MB, **Snyder M**, Myers RM, Weng Z. Integrated analysis of experimental data sets reveals many novel promoters in 1% of the human genome. *Genome Res.* 2007;17: 720-31.
19. Rozowsky JS, Newburger D, Sayward F, Wu J, Jordan G, Korbel JO, Nagalakshmi U, Yang J, Zheng D, Guigo R, Gingeras TR, Weissman S, Miller P, **Snyder M**, Gerstein MB. The DART classification of unannotated transcription within the ENCODE regions: associating transcription with known and novel loci. *Genome Res.* 2007;17: 732-45.
20. Zhang ZD, Paccanaro A, Fu Y, Weissman S, Weng Z, Chang J, **Snyder M**, Gerstein MB. Statistical analysis of the genomic distribution and correlation of regulatory elements in the ENCODE regions. *Genome Res.* 2007;17: 787-97.
21. Zheng D, Frankish A, Baertsch R, Kapranov P, Reymond A, Choo SW, Lu Y, Denoeud F, Antonarakis SE, **Snyder M**, Ruan Y, Wei CL, Gingeras TR, Guigo R, Harrow J, Gerstein MB. Pseudogenes in the ENCODE regions: consensus annotation, analysis of transcription, and evolution. *Genome Res.* 2007;17: 839-51.
22. Washietl S, Pedersen JS, Korbel JO, Stocsits C, Gruber AR, Hackermuller J, Hertel J, Lindemeyer M, Reiche K, Tanzer A, Ucla C, Wyss C, Antonarakis SE, Denoeud F, Lagarde J, Drenkow J, Kapranov P, Gingeras TR, Guigo R, **Snyder M**, Gerstein MB, Reymond A, Hofacker IL, Stadler PF. Structured RNAs in the ENCODE selected regions of the human genome. *Genome Res.* 2007;17: 852-64.
23. Bhinge AA, Kim J, Euskirchen GM, **Snyder M**, Iyer VR. Mapping the chromosomal targets of STAT1 by Sequence Tag Analysis of Genomic Enrichment (STAGE). *Genome Res.* 2007;17: 910-6.
24. Borneman AR, Zhang ZD, Rozowsky J, Seringhaus MR, Gerstein M, **Snyder M**. Transcription factor binding site identification in yeast: a comparison of high-density oligonucleotide and PCR-based microarray platforms. *Funct Integr Genomics.* 2007;7: 335-45.
25. Borneman AR, Gianoulis TA, Zhang ZD, Yu H, Rozowsky J, Seringhaus MR, Wang LY, Gerstein M, **Snyder M**. Divergence of transcription factor binding sites across related yeast species. *Science.* 2007;317: 815-19.
26. Devgan G, **Snyder M**. Kinase substrate identification using yeast protein micorarrays. In: Functional Microarrays in Drug Discovery. P Predki, ed. 2007. CRC Press 351-360.
27. Popescu SC, Popescu GV, Bachan S, Zhang Z, Seay M, Gerstein M, **Snyder M**, Dinesh-

- Kumar SP. Differential binding of calmodulin related proteins to their targets revealed using high density Arabidopsis protein microarrays. *Proc Natl Acad Sci USA*. 2007;104: 4730-5.
28. Rozowsky J, Wu J, Lian Z, Nagalakshmi U, Korbel JO, Kapranov PD, Zheng D, Dyke S, Newburger P, Miller P, Gingeras T, Weissman S, Gerstein M, **Snyder M**. Novel transcribed regions in the human genome. *Cold Spring Harb Symp Quant Biol*. 2007;71: 111-116.
 29. Euskirchen GM, Rozowsky J, Wei CL, Lee WH, Zhang ZD, Hartman S, Emanuelsson O, Stolc V, Weissman S, Gerstein M, Ruan Y, **Snyder M**. Mapping of transcription factor binding regions in mammalian cells by ChIP: comparison of array- and sequencing-based technologies. *Genome Res*. 2007;17: 898-909.
 30. Smith MG, Gianoulis TA, Pukatzki S, Mekalanos J, Ornston LN, Gerstein M, **Snyder M**. New insights into *Acinetobacter baumannii* pathogenesis revealed by high-density pyrosequencing and transposon mutagenesis. *Genes Dev*. 2007;21: 601-14.
 31. Hall DA, Ptacek J, **Snyder M**. Protein microarray technology. *Mech Aging Dev*. 2007;128: 161-7.
 32. Emanuelsson O, Nagalakshmi U, Zheng D, Rozowsky JS, Urban AE, Du J, Lian Z, Stolc V, Weissman S, **Snyder M**, Gerstein M. Assessing the performance of different high-density tiling microarray strategies for mapping transcribed regions of the human genome. *Genome Res*. 2007;17: 886-97.
 33. Yu H, Nguyen K, Royce T, Qian J, Nelson K, **Snyder M**, Gerstein M. Positional artifacts in microarrays: experimental verification and construction of COP, an automated detection tool. *Nucleic Acids Res*. 2007;35.e8.
 34. **Snyder M**. Yeast is still the beast. *GSA Newsletter*. Fall 2006.
 35. Dewan A, Liu M, Hartman S, Zhang SS, Liu DT, Zhao C, Tam PO, Chan WM, Lam DS, **Snyder M**, Barnstable C, Pang CP, Hoh J. HTRA1 promoter polymorphism in wet age-related macular degeneration. *Science*. 2006;314: 989-92.
 36. Wang LY, **Snyder M**, Gerstein M. BoCaTFBS: a boosted cascade learner to refine the binding sites suggested by ChIP-chip experiments. *Genome Biol*. 2006;7:R102.
 37. Zhu X, Gerstein M, **Snyder M**. ProCAT: a data analysis approach for protein microarrays. *Genome Biol*. 2006;7: R110.
 38. Hudson ME, **Snyder M**. High-throughput methods of regulatory element discovery. *Biotechniques*. 2006;41: 673-677.
 39. Du J, Rozowsky JS, Korbel JO, Zhang ZD, Royce TE, Schultz MH, **Snyder M**, Gerstein M. A supervised hidden markov model framework for efficiently segmenting tiling array data in transcriptional and ChIP-chip experiments: systematically incorporating validated biological knowledge. *Bioinformatics*. 2006;22: 3016-24.
 40. Seringhaus M, Paccanaro A, Borneman A, **Snyder M**, Gerstein M. Predicting essential genes in fungal genomes. *Genome Res*. 2006;16: 1126-35.
 41. Ptacek J, **Snyder M**. Charging it up: global analysis of protein phosphorylation. *Trends Genet*. 2006;22: 545-54

42. Srikantha T, Borneman AR, Daniels KJ, Pujol C, Wu W, Seringhaus MR, Gerstein M, Yi S, **Snyder M**, Soll DR. TOS9 regulates white-opaque switching in *Candida albicans*. *Eukaryot Cell*. 2006;5: 1674-87.
43. Kastenmayer JP,* Ni L,* Chu A, Kitchen LE, Au WC, Yang H, Carter CD, Wheeler D, Davis RW, Boeke JD, **Snyder MA**, Basrai MA. Functional genomics of genes with small open reading frames (sORFs) in *S. cerevisiae*. *Genome Res*. 2006;16: 365-73.
44. Seringhaus M, Kumar A, Hartigan J, **Snyder M**, Gerstein M. Genomic analysis of insertion behavior and target specificity of mini-Tn7 and Tn3 transposons in *Saccharomyces cerevisiae*. *Nucleic Acids Res*. 2006;34: e57.
45. Kung LA, **Snyder M**. Proteome chips for whole-organism assays. *Nat Rev Mol Cell Biol*. 2006;7: 617-22.
46. Sopko R, Huang D, Preston N, Chua G, Papp B, Kafadar K, **Snyder M**, Oliver SG, Cyert M, Hughes TR, Boone C, Andrews B. Mapping pathways and phenotypes by systematic gene overexpression. *Mol Cell*. 2006;21: 319-30.
47. **Snyder M**, Gelperin D. Community resources for *Saccharomyces cerevisiae*. 2006, Jan/Feb 9.
48. Lu J, Hou R, Booth CJ, Yang S-H, **Snyder M**. Defined culture of human embryonic stem cells. *Proc Natl Acad Sci USA*. 2006;103: 5688-93.
49. Royce TE, Rozowsky JS, Luscombe NM, Emanuelsson O, Yu H, Zhu X, **Snyder M**, Gerstein M. Extrapolating traditional DNA microarray statistics to the tiling and protein microarray technologies. *Meth Enzymol*. 2006;411: 282-311.
50. Ho S, Jona G, Johnston M, **Snyder M**. Linking DNA-binding proteins to their recognition sequences using protein microarrays. *Proc Natl Acad Sci USA*. 2006;103: 9940-45.
51. Smith MG, **Snyder M**. Yeast as a model for human disease. *Current Protocols in Human Genetics*. 2006;15.6.1-15.6.8.
52. Urban AE, Korbel J, Selzer R, Popescu GV, Richmond T, Cubells JF, Green R, Emanuel BS, Gerstein M, Weissman SM, **Snyder M**. High resolution mapping of DNA copy alterations using high density tiling oligonucleotide arrays. *Proc Natl Acad Sci USA*. 2006;103: 4534-9.
53. Borneman R, Leigh-Bell J, Yu H, Bertone P, Gerstein M, **Snyder M**. Target hub proteins serve as master regulators of the complex transcriptional network controlling yeast pseudohyphal growth. *Genes Devel*. 2006;20: 435-448.
54. Zhu H, Hu S, Jona G, Zhu X, Kreiswirth N, Willey BM, Mazzulli T, Liu G, Song Q, Chen P, Cameron M, Tyler A, Wang J, Wen J, Chen W, Compton S, **Snyder M**. SARS diagnostics using a coronavirus protein chip. *Proc Nat Acad Sci USA*. 2006;103: 4011-16.
55. Bertone P, Trifonov V, Rozowsky JS, Schubert F, Emanuelsson O, Karro J, Kao M-Y, **Snyder M**, Gerstein M. Design optimization methods for genomic DNA tiling arrays. *Genome Res*. 2006;16: 271-81.
56. Stolc V, Li L, Wang X, Li X, Su N, Tongprasit W, Han B, Xue Y, Li J, **Snyder M**, Gerstein M, Wang J, Deng XW. A pilot study of transcription unit analysis in rice using oligonucleotide tiling-path microarray. *Plant Mol Biol*. 2005;59: 137-49.

57. Mah AS, Elia AE, Devgan G, Ptacek J, Schutkowski M, **Snyder M**, Yaffe MB, Deshaies RJ. Substrate specificity analysis of protein kinase complex Dbf2-Mob1 by peptide library and proteome array screening. *BMC Biochem.* 2005;6: 22.
58. Hartman SE, Bertone P, Nath A, Royce TE, Gerstein M, Weissman S, **Snyder M**. Global changes in STAT target selection and transcription regulation upon interferon treatments. *Genes Devel.* 2005;19: 2953-68.
59. Royce TE, Rozowsky JS, Bertone P, Samantac M, Stolc V, Weissman S, **Snyder M**, Gerstein M. Issues in the analysis of oligonucleotide tiling microarrays for transcript mapping. *Trends Genet.* 2005;21: 466-75.
60. Bertone P, **Snyder M**. Advances in functional protein microarrays. *FEBS J.* 2005;272: 5400-5411.
61. Ptacek J, Devgan G, Michaud G, Zhu H, Zhu X, Fasolo J, Guo H, Jona G, Breitkreutz A, Sopko R, Lee S, McCartney RR, Schmidt MC, Rachidi N, Stark MJR, Stern DF, Tyers M, de Virgilio C, Andrews B, Gerstein M, Schweitzer B, Predki P, **Snyder M**. Global analysis of protein phosphorylation in yeast. *Nature.* 2005;438: 679-84.
62. Gelperin DM, White MA, Wilkinson ML, Kon Y, Li A, Kung LA, Wise KJ, Lopez-Hoyo N, Jiang L, Piccirillo S, Yu H, Gerstein M, Dumont ME, Phizicky EM, **Snyder M**,* Grayhack EJ.* Biochemical and genetic analysis of the yeast proteome with a movable ORF collection. *Genes Devel.* 2005;19: 2816-26. *Co-Senior Authors
63. Pot I, Knockleby J, Aneliunas V, Nguyen T, Ah-Kye S, Liszt G, **Snyder M**, Hieter P, Vogel J. Spindle checkpoint maintenance requires Amel1 and Okp1. *Cell Cycle.* 2005;4: 1448-56.
64. Oh SW, Moon JD, Lim HJ, Park SY, Kim T, Park J, Han MH, **Snyder M**, Choi EY. Calixarene derivative as a tool for highly sensitive detection and oriented immobilization of proteins in a microarray format through noncovalent molecular interaction. *FASEB J.* 2005;19: 1335-7.
65. Bertone P, **Snyder M**. Prospects and challenges in proteomics. *Plant Physiol.* 2005;138: 560-2.
66. Bertone P, Gerstein M, **Snyder M**. Applications of DNA tiling arrays to experimental genome annotation and regulatory pathway discovery. *Chromosome Res.* 2005;13: 259-74.
67. Martone R, **Snyder M**. Mapping transcription factor binding sites using chIP Chip - general considerations. *DNA Microarrays.* 2005, in press.
68. Rinn JL, **Snyder M**. Sexual dimorphism in mammalian gene expression. *Trends Genet.* 2005;21: 298-305.
69. Smith MG, Jona G, Ptacek J, Devgan G, Zhu H, Zhu X, **Snyder M**. Global analysis of protein function using protein microarrays. *Mechanisms Ageing Devel.* 2005;126: 171-175.
70. ENCODE Project Consortium. The ENCODE (EnCyclopedia Of DNA Elements) Project. *Science.* 2004;306: 636-640.
71. Bertone P, Stolc V, Royce TE, Rozowsky JS, Urban AE, Zhu X, Rinn JL, Tongprasit W, Samanta M, Weissman S, Gerstein M, **Snyder M**. Global identification of human transcribed sequences with genome tiling arrays. *Science.* 2004;306: 2242-6.

72. Mukherjee S, Berger MF, Jona G, Wang XS, Muzzey D, **Snyder M**, Young RA, Bulyk ML. Rapid analysis of the DNA-binding specificities of transcription factors with DNA microarrays. *Nat Genet.* 2004;36: 1331-9.
73. White EJ, Emanuelsson O, Scalzo D, Royce T, Kosak S, Weissman S, Gerstein M, Groudine M, **Snyder M**, Schubeler D. High resolution DNA replication analysis of human chromosome 22 reveals cell type specific differences in DNA replication timing. *Proc Natl Acad Sci USA.* 2004;101: 17771-6.
74. Kumar A, Seringhaus M, Biery MC, Sarnovsky RJ, Umansky L, Piccirillo S, Matson S, Heidtman M, Cheung KH, Dobry CJ, Gerstein M, Craig NL, **Snyder M**. Large-scale mutagenesis of the yeast genome using a Tn7-derived multipurpose transposon. *Genome Res.* 2004;14: 1975-86.
75. Luscombe NM, Babu MM, Yu H, **Snyder M**, Teichmann SA, Gerstein M. Genome analysis of regulatory network dynamics reveals large topological changes. *Nature.* 2004;431: 308-12.
76. Hall DA, Zhu H, Royce T, Gerstein M, **Snyder M**. Regulation of gene expression by a metabolic enzyme. *Science.* 2004;306: 482-484.
77. Berman P, Bertone P, DasGupta B, Gerstein M, Ming-Yang Kao M-Y, **Snyder M**. Fast optimal genome tiling with applications to microarray design and homology search. *J. Computational Biol.* 2004;11: 766-785.
78. Euskirchen G, **Snyder M**. A plethora of sites. *Nat Genet.* 2004;36: 325-326.
79. Rinn JL, Rozowsky JS, Laurenz IJ, Petersen PH, Zou K, Zhong W, Gerstein M, **Snyder M**. Major molecular differences between mammalian sexes are involved in drug metabolism and renal function. *Dev Cell.* 2004;6: 791-800.
80. Euskirchen G, Royce TE, Bertone P, Martone R, Rinn JL, Nelson FK, Sayward F, Luscombe NM, Miller P, Gerstein M, Weissman S, **Snyder M**. CREB binds to multiple loci on chromosome 22. *Mol Cell Biol.* 2004;24: 3804-3814.
81. Smith M, des Etages S, **Snyder M**. Microbial synergy via an ethanol triggered pathway. *Mol Cell Biol.* 2004;24: 3874-3884.
82. Bidlingmaier S, **Snyder M**. Regulation of polarized growth initiation and termination cycles by the polarisome and Cdc42 regulators. *J Cell Biol.* 2004;164: 201-18. (Featured on cover).
83. Huang J, Zhu H, Haggarty SJ, Spring DR, **Snyder M**, Schreiber S. Finding new components of the TOR signaling pathway using chemical genetics and proteome chips. *Proc Natl Acad Sci USA.* 2004;101: 16594-9.
84. Lian Z, Euskirchen G, Rinn J, Martone R, Bertone P, Hartman S, Royce T, Nelson K, Sayward F, Luscombe N, Yang J, Li JL, Miller P, Urban AE, Gerstein M, Weissman S, **Snyder M**. Identification of novel functional elements in the human genome. *Cold Spring Harb Symp Quant Biol.* 2003;68: 317-322.
85. Jansen R, Yu H, Greenbaum D, Kluger Y, Krogan NJ, Chung S, Emili A, **Snyder M**, Greenblatt JF, Gerstein M. A Bayesian networks approach for predicting protein-protein interactions from genomic data. *Science.* 2003;302: 449-53.

86. Michaud GA, Salcius M, Zhou F, Bangham R, Bonin J, Guo H, **Snyder M**, Predki PF, Schweitzer BI. Analyzing antibody specificity with whole proteome microarrays. *Nat Biotechnol.* 2003;21: 1509-12.
87. Kafadar KA, Zhu H, **Snyder M**, Cyert M. Negative regulation of calcineurin signaling by Hrr25p, a yeast homolog of casein kinase I. *Genes Devel.* 2003;17: 2698-708.
88. Martone R, Euskirchen G, Bertone P, Hartman S, Royce TE, Luscombe NL, Rinn JL, Nelson FK, Miller P, Gerstein M, Weissman S, **Snyder M**. Distribution of NF- κ B binding sites across human chromosome 22. *Proc Natl Acad Sci USA.* 2003;100: 12247-12252.
89. Schweitzer B, Predki P, **Snyder M**. Microarrays to characterize protein interactions on a whole-proteome scale. *Proteomics.* 2003;3: 2190-9.
90. Ball J, Schweitzer B, Predki P, **Snyder M**. Development and applications of functional protein arrays. M Schena ed. Jones and Bartlett Publishers, Boston. Protein Microarrays. 2003. p421-440.
91. Phizicky E, Bastiaens PI, Zhu H, **Snyder M**, Fields S. Protein analysis on a proteomic scale. *Nature.* 2003;422: 208-15.
92. Hanrahan J, **Snyder M**. Cytoskeletal activation of a checkpoint kinase. *Mol Cell.* 2003;12: 663-73. (Featured on cover)
93. Jona G, **Snyder M**. Recent developments in analytical and functional protein microarrays. *Curr Opin Mol Ther.* 2003;3: 271-7.
94. Santos B, **Snyder M**. Protein targeting during cell differentiation: polarized localization of Fus1p during mating depends on Chs5p in *Saccharomyces cerevisiae*. *Eucaryotic Cell.* 2003;2: 821-825.
95. Reboul J, Vaglio P, Rual JF, Lamesch P, Martinez M, Armstrong CM, Li S, Jacotot L, Bertin N, Janky R, Moore T, Hudson JR Jr, Hartley JL, Brasch MA, Vandenhaute J, Boulton S, Endress GA, Jenna S, Chevet E, Papasotiropoulos V, Tolia PP, Ptacek J, **Snyder M**, Huang R, Chance MR, Lee H, Doucette-Stamm L, Hill DE, Vidal M. *C. elegans* ORFeome version 1.1: experimental verification of the genome annotation and resource for proteome-scale protein expression. *Nat Genet.* 2003;34: 35-41.
96. Luscombe NM Royce TE, Bertone P, Echols N, Horak CE, Chang JT, **Snyder M**, Gerstein M. Express yourself: a modular platform for processing and visualizing microarray data. *Nucleic Acids Res.* 2003;31: 3477-82.
97. Zhu H, Bilgin M, **Snyder M**. Proteomics. *Ann Rev Biochem.* 2003;72: 783-812.
98. **Snyder M**, Gerstein M. Defining genes in the genomics era. *Science.* 2003;300: 258-260.
99. Casamayor A, **Snyder M**. Molecular dissection of a yeast septin: distinct domains are required for septin interaction, localization and function. *Mol Cell Biol.* 2003;23: 2762-2777.
100. Zhu H, **Snyder M**. Protein chip technology. *Curr Opin Chem Biol.* 2003;7: 55-63.
101. Rinn JR, Euskirchen G, Bertone P, Martone R, Luscombe NM, Hartman S, Harrison PM, Nelson FN, Miller P, Gerstein M, Weissman S, **Snyder M**. The transcriptional activity of human chromosome 22. *Genes Devel.* 2003;17: 529-40.

102. Cheung KH, White K, Hager J, Gerstein M, Reinke V, Nelson K, Masiar P, Srivastava R, Li Y, Li J, Zhao H, Li J, Allison DB, **Snyder M**, Miller P, Williams K. YMD: a microarray database for large-scale gene expression analysis. *Proc AMLA Symp.* 2002: 140-4.
103. Michaud G, **Snyder M**. Proteomic approaches for global analysis of proteins. *Biotechniques.* 2002;33: 1308-18.
104. Horak CE, Luscombe NM, Qian J, Piccirillo S, Gerstein M, **Snyder M**. Complex transcriptional circuitry at the G1/S transition in *Saccharomyces cerevisiae*. *Genes Dev.* 2002;16: 3017-33.
105. Coelho PSR, Bryan AC, Kumar A, Shadel GS, **Snyder M**. A novel mitochondrial protein, Tar1p, is encoded on the antisense strand of the nuclear 25S rDNA. *Genes Devel.* 2002;16: 2755-2760.
106. Vallier L, Segall J, **Snyder M**. The alpha-factor receptor C-terminus is important for mating projection formation and orientation in *Saccharomyces cerevisiae*. *Cell Motil Cytoskeleton.* 2002;53: 251-266.
107. Giaever G, Chu AM, Ni L, Connelly C, Riles L, Veronneau S, Dow S, Lucau-Danila A, Anderson K, Andre B, Arkin AP, Astromoff A, El-Bakkoury M, Bangham R, Benito R, Brachat S, Campanaro S, Curtiss M, Davis K, Deutschbauer A, Entian KD, Flaherty P, Foury F, Garfinkel DJ, Gerstein M, Gotte D, Guldener U, Hegemann JH, Hempel S, Herman Z, Jaramillo DF, Kelly DE, Kelly SL, Kotter P, LaBonte D, Lamb DC, Lan N, Liang H, Liao H, Liu L, Luo C, Lussier M, Mao R, Menard P, Ooi SL, Revuelt JL, Roberts CJ, Rose M, Ross-Macdonald P, Scherensm B, Schimmack G, Shafer B, Shoemaker DD, Sookhai-Mahadeo S, Storms RK, Strathern JN, Valle G, Voet M, Volckaert G, Wang CY, Ward TR, Wilhelmy J, Winzeler EA, Yang Y, Yen G, Youngman E, Yu K, Bussey H, Boeke JD, **Snyder M**, Philippsen P, Davis RW, Johnston M. Functional profiling of the *Saccharomyces cerevisiae* genome. *Nature.* 2002;418: 387-91.
108. Horak CE, **Snyder M**. Global analysis of gene expression in yeast. *Funct Integr Genomics.* 2002;2: 171-80.
109. **Snyder M**, Kumar A. Yeast genomics: past, present, and future promise. *Funct Integr Genomics.* 2002;2: 135-137.
110. Bidlingmaier S, **Snyder M**. Carbohydrate analysis prepares to enter the "Omics" era. *Chem Biol.* 2002;9: 400-1.
111. Horak CE, Mahajan MC, Luscombe NM, Gerstein M, Weissman SM, **Snyder M**. GATA-1 binding sites mapped in the β -globin locus by using mammalian ChIp-chip analysis. *Proc Natl Acad Sci USA.* 2002;99: 2924-29.
112. Kusch J, Meyer A, **Snyder M**, Barral Y. Proper positioning of the spindle and the cleavage plane relative to each other involves microtubule capture by the cleavage apparatus in yeast. *Genes Dev.* 2002;16: 1627-1639
113. Zhu H, **Snyder M**. "OMIC" approaches to unraveling cell signaling. *Curr Opin Cell Biol.* 2002;14: 173-179.
114. Casamayor A, **Snyder M**. Bud site selection and cell polarity in budding yeast. *Curr Opin Microbiol.* 2002;5: 179-86.

115. Harrison PM, Kumar A, **Snyder M**, Gerstein MB. A question of size: the eukaryotic proteome and the problems in defining it. *Nucl Acids Res.* 2002;30: 1083-90.
116. Kumar A, Cheung K-H, Marenco L, Tosches N, Masiar P, Liu Y, Miller P, **Snyder M**. The TRIPLES database of yeast protein function. In: *Analyzing Gene Expression* (ed., Lorkowski, S. and Cullen, P., et al.) Wiley-VCH Verlag GmbH, Weinheim, Germany. 2002, in press.
117. Harrison P, Kumar A, Lan N, Echols N, **Snyder M**, Gerstein MB. A small reservoir of disabled ORFs in the *Saccharomyces cerevisiae* genome and its implication for the dynamics of proteome evolution. *J Mol Biol.* 2002;316: 409-19.
118. Kumar A, Cheung K-H, Tosches N, Masiar P, Liu Y, Miller P, **Snyder M**. The TRIPLES database: a community resource for yeast molecular biology. *Nucl Acids Res.* 2002;30: 73-75.
119. Kumar A, **Snyder M**. Protein complexes take the bait. *Nature.* 2002;415: 123-124.
120. Kumar A, Agarwal S, Heyman JA, Matson S, Heidtman M, Piccirillo S, Umansky L, Drawid A, Jansen R, Liu Y, Cheung K-H, Miller P, Gerstein MB, Roeder GS, **Snyder M**. Subcellular localization of the yeast proteome. *Genes Dev.* 2002;16: 707-719.
121. Kumar A, Vidan S, **Snyder M**. Insertional mutagenesis: transposon-insertion libraries as mutagens in yeast. *Methods Enzymol.* 2002;350: 219-229.
122. Horak C, **Snyder M**. ChIP chip: a genomic approach for identifying transcription factor binding sites. *Methods Enzymol.* 2002;350: 469-484.
123. Bidlingmaier S, **Snyder M**. Large-scale identification of genes important for apical growth in *Saccharomyces cerevisiae* by direct allele replacement technology (DART) screening. *Funct Integr Genomics.* 2002;1: 345-356.
124. Kumar A, Harrison PS, Cheung K-H, Lan N, Echols N, Bertone P, Miller P, Gerstein MB, **Snyder M**. An integrated approach for finding overlooked genes in yeast. *Nat Biotech.* 2002;20: 58-63. (Featured in *Nat Biotech News and Views*).
125. Cheung KH, Deshpande AM, Tosches N, Nath S, Agrawal A, Miller P, Kumar A, **Snyder M**. A metadata framework for interoperating heterogeneous genome data using XML. *Proc AMIA Symp.* 2001; 110-4.
126. Vidan S, **Snyder M**. Making drug addicts out of yeast. *Nature Biotech.* 2001;19: 1022-1023.
127. Vogel J, Drapkin B, Oomen J, Beach D, Bloom K, **Snyder M**. Phosphorylation of gamma-tubulin regulates microtubule organization in budding yeast. *Dev Cell.* 2001;1: 621-631.
128. Zhu H, **Snyder M**. Biochemical assays in a chip format. *Current Drug Discovery.* 2001;31-34.
129. Zhu H, Bilgin M, Bangham R, Hall D, Casamayor A, Bertone P, Lan N, Jansen R, Bidlingmaier S, Houfek T, Mitchell T, Miller P, Dean DA, Gerstein M, **Snyder M**. Global analysis of protein activities using proteome chips. *Science.* 2001;293: 2101-2105. (Featured in many journals, websites and newspapers.)
130. Sheu Y-J, **Snyder M**. Control of cell polarity and shape in *S. cerevisiae*. Howard and N.

Gow eds. Springer, Heidelberg. Mycota Vol VIII: The Biology of the Fungal Cell. 2001 p. 19-53.

131. Cheung KH, Hager J, White K, Williams K, Nelson K, **Snyder M**, Li Y, Miller P. MAC: a Web interface for convoluting data from plates to microarrays. 2001, in press.
132. Ni L, **Snyder M**. A genomic study of the bipolar bud site selection pattern in *Saccharomyces cerevisiae*. *Mol Biol Cell*. 2001;12: 2147-70.
133. Kumar A, **Snyder M**. Emerging technologies in yeast genomics. *Nature Rev Genet*. 2001;2: 302-312.
134. Bidlingmaier S, Weiss EL, Siedel C, Drubin DG, **Snyder M**. The Cbk1 pathway is important for polarized cell growth and cell separation in *Saccharomyces cerevisiae*. *Mol Cell Biol*. 2001;21: 2449-62.
135. Iyer VI,* Horak CA,* Scafe CS, Botstein D, **Snyder M**, Brown PO. Genomic binding distribution of the yeast cell-cycle transcription factors SBF and MBF. *Nature*. 2001;409: 533-538.
136. Vidan S, **Snyder M**. Large-scale mutagenesis: yeast genetics in the genome era. *Curr Opin Biotech*. 2001;12: 28-34.
137. Zhu H, **Snyder M**. Protein arrays and microarrays. *Curr Opin Chem Biol*. 2001;5: 40-45.
138. Erdman S, **Snyder M**. A filamentous growth response mediated by the yeast mating pathway. *Genetics*. 2001;159: 919-928.
139. Cheung K, Miller P, Sherman A, Weston S, Stratmann E, Schultz M, **Snyder M**, Kumar A. Graphically-enabled integration of bioinformatics tools allowing parallel execution. *Proc. AMIA Symp*. 2000;20: 141-5.
140. Zhu H, Klemic JF, Chang S, Bertone P, Klemic KG, Smith D, Gerstein M, Reed MA, **Snyder M**. Analysis of yeast protein kinases using protein chips. *Nat Genet*. 2000;26: 283-289.
141. Bouquin N, Barral Y, Courbeyrette R, Blondel M, **Snyder M**, Mann C. Regulation of cytokinesis by the Elm1 protein kinase in *Saccharomyces cerevisiae*. *J Cell Sci*. 2000;113: 1435-45.
142. Vogel J, **Snyder M**. The carboxy terminus of Tub4p is required for gamma-tubulin function in budding yeast. *J Cell Sci*. 2000;113: 3871-82.
143. Coehlo PSR, Kumar A, **Snyder M**. Genome-wide mutant collections: toolboxes for functional genomics. *Curr Opin Microbiol*. 2000;3: 309-315.
144. Deng Y, Lee JP, Tianasoa-Ramamonjy M, Snyder M, Des Etages SA, Kanada D, **Snyder MP**, Turner CJ. New Antimicrobial flavanones from *Physena madagascariensis*. *J Natural Products*. 2000;63: 1082-9.
145. Manning BD, **Snyder M**. Drivers and passengers wanted: kinesin associated proteins. *Trends Cell Biol*. 2000;10: 281-289.
146. Barrett JG, Manning BD, **Snyder M**. The Kar3p kinesin-related protein forms a novel

- heterodimer structure with its associated protein Cik1p. *Mol Biol Cell*. 2000;11: 2373-2385.
147. Barral Y, Mermall V, Mooseker MS, **Snyder M**. The septins form a barrier necessary for the maintenance of cell polarity in yeast. *Molecular Cell*. 2000;5: 841-851.
148. Kumar A, **Snyder M**. Genome-wide transposon mutagenesis in yeast. In: Current Protocols in Molecular Biology, 2000. (ed, Ausubel, F.M., Brent, R., Kingston, R.E., Moore, D., Seidman, J.G., Smith, J.A., and Struhl, K.) Wiley and Sons, N.Y., N.Y. Unit 13.3.
149. Sheu Y-J, Barral Y, **Snyder M**. Polarized growth controls cell shape and bud site selection in *Saccharomyces cerevisiae*. *Mol Cell Biol*. 2000;14: 5235-5247.
150. Cheung KH, Kumar A, **Snyder M**, Miller P. An integrated web interface for large-scale characterization of sequence data. *Funct Integr Genomics*. 2000;1: 70-75.
151. Zhan X, Lee M, Abenes G, Von Reis I, Kittinunvorakoon C, Ross-Macdonald P, **Snyder M**, Liu F. Mutagenesis of murine cytomegalovirus using a Tn3-based transposon. *Virology*. 2000;266: 264-74.
152. Santos B, **Snyder M**. Sbe2p and Sbe22p, two homologous Golgi proteins involved in yeast cell wall formation. *Mol Biol Cell*. 2000;11: 435-452.
153. Kumar A, des Etages SA, Coelho PSR, Roeder GS, **Snyder M**. High-throughput methods for the large-scale analysis of gene function by transposon tagging. *Methods Enzymol*. 2000;328: 550-74.
154. Kumar A, Cheung K-H, Ross-Macdonald P, Coelho PSR, Miller P, **Snyder M**. TRIPLES: a database of gene function in *S. cerevisiae*. *Nuc Acids Res*. 2000;28: 81-84.
155. Mattagajasingh SN, Huang SC, Hartenstein JS, **Snyder M**, Marchesi VT, Benz EJ. A nonerythroid isoform of protein 4.1R interacts with the nuclear mitotic apparatus (NuMA) protein. *J Cell Biol*. 1999;145: 29-43.
156. des Etages SA, Kumar A, **Snyder M**. Transposons as tools. In: Encyclopedia of Genetics, 1999. S Brenner and JH Miller, eds. Academic Press, San Diego. p. 2034-2040.
157. Vogel J, **Snyder M**. The centrosome in cell replication and development. *Curr Topics Devel*. 1999;49: 75-104.
158. Ross-Macdonald P, Coelho PSR, Roemer T, Agarwal S, Kumar A, Jansen R, Cheung K-H, Sheehan A, Symoniatis D, Umansky L, Heitman M, Nelson FK, Iwasaki H, Hager K, Gerstein M, Miller P, Roeder GS, **Snyder M**. Large-scale analysis of the yeast genome by transposon tagging and gene disruption. *Nature*. 1999;402: 413-418. (Featured in News and Views.)
159. Winzeler EA, Astromoff A, Liang H, Shoemaker DD, Anderson K, Bangham R, Boeke JD, Bussey H, Connelly C, Davis K, Dietrich F, Foury F, Friend S, Gentalen E, Giaever G, Jones T, Laub M, Liao H, Lockhart D, Lussier M, Ross-Macdonald P, Menard P, Mittmann M, Pai C, Philippsen P, Rebischung C, Riles L, Rine J, Roberts C, **Snyder M**, Stoms RK, Veronneau S, Ward T, Whelan S, Yen G, Yu K, Johnston M, Davis RW. Functional characterization of the *Saccharomyces cerevisiae* genome by comprehensive and precise gene deletion and massively parallel analysis. *Science*. 1999;285: 901-906.
160. Manning BD, Barrett J, Wallace J, Granok H, **Snyder M**. Differential regulation of the

- Kar3p kinesin-related protein by two associated proteins, Vik1p and Cik1p. *J Cell Biol.* 1999;144: 1219-1233.
161. Hong SK, Han SB, **Snyder M**, Choi EY. *SHC1*, a high pH inducible gene required for growth at alkaline pH in *Saccharomyces cerevisiae*. *Biochem Biophys Res Comm.* 1999;255: 116-122.
162. Barral Y, Parra M, Bidlingmaier S, **Snyder M**. Nim1-related kinases coordinate cell cycle progression with organization of the peripheral cytoskeleton. *Genes Dev.* 1999;13: 176-187. (Featured on Cover.)
163. Ross-MacDonald P, Sheehan A, Friddle C, Roeder GS, **Snyder M**. Transposon mutagenesis for the analysis of protein production, function and localization. *Methods Enzymol.* 1999;303: 512-532.
164. Madden K, **Snyder M**. Cell polarity and morphogenesis in budding yeast. *Ann Rev Microbiol.* 1998;52: 687-744.
165. Sheu Y-J, Santos B, Fortin N, Costigan C, **Snyder M**. Spa2p interacts with cell polarity proteins and signaling components involved in yeast cell morphogenesis. *Mol Cell Biol.* 1998;18: 4053-4069.
166. Roemer T, Vallier L, Sheu Y-J, **Snyder M**. The yeast Sph1 protein is important for cell morphogenesis in yeast. *J Cell Science.* 1998;111: 479-494.
167. Erdman S, Lin L, Malczynski M, **Snyder M**. Pheromone-regulated genes required for yeast mating differentiation. *J Cell Biol.* 1998;140: 461-483.
168. Ross-MacDonald P, Sheehan A, Friddle C, Roeder GS, **Snyder M**. Transposon tagging: a novel system for monitoring protein production, function and localization. In: *Methods in Microbiology: Yeast Gene Analysis*, 1998. M.F. Tuite & A.J.P. Brown, eds, Academic Press, London, Vol 26, 161-179.
169. Costigan C, **Snyder M**. Cell polarity in the budding yeast, *Saccharomyces cerevisiae*. *Adv Mol Cell Biol.* 1998;26: 1-66.
170. Manning BD, Padmanabha R, **Snyder M**. The Rho-GEF Rom2p localizes to sites of polarized cell growth and participates in cytoskeletal functions in *Saccharomyces cerevisiae*. *Mol Biol Cell.* 1997;8: 1829-1844.
171. Semenov MV, **Snyder M**. Human dishevelled genes comprise a DHR-containing multigene family. *Genomics.* 1997;42: 302-310.
172. Madden K, Sheu Y-J, Baetz K, Andrews B, **Snyder M**. SBF cell cycle regulator as a target of yeast PKC-MAP kinase pathway. *Science.* 1997;275: 1781-1784.
173. Santos B, **Snyder M**. Targetting of chitin synthase 3 to polarized growth sites in yeast requires Chs5p and Myo2p. *J Cell Biol.* 1997;136: 95-110.
174. Ross-MacDonald P, Sheehan A, Roeder GS, **Snyder M**. A multipurpose transposon system for analyzing protein production, localization and function in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci USA.* 1997;94: 190-195.
175. Roemer T, Vallier L, **Snyder M**. Selection of polarized growth sites in yeast. *Trends Cell*

Biol. 1996;6: 434-441.

176. Roemer T, Madden K, Chang J, **Snyder M**. Selection of axial growth sites in yeast requires Axl2p, a novel plasma membrane glycoprotein. *Genes Devel.* 1996;10: 777-793.
177. Chen CR, Malik M, **Snyder M**, Drlica K. DNA gyrase and topoisomerase IV on the bacterial chromosome: quinolone-induced DNA cleavage. *J Mol Biol.* 1996;258: 627-637.
178. Burns N, Ross-Macdonald P, Roeder GS, **Snyder M**. Generation, screening and analysis of *lacZ* fusions in yeast. In: *Microbial Genome Methods*, 1996. K. Adolph, ed. CRC Press, New York p. 61-79.
179. Friedman-Einat M, Einat P, **Snyder M**, Ruddle F. Target gene identification: target specific transcriptional activation by three murine homeodomain/VP16 hybrid proteins in *Saccharomyces cerevisiae*. *J Exp Zoology.* 1996;274: 145-156.
180. Ross-Macdonald P, Burns N, Malczynski M, Sheehan A, Roeder GS, **Snyder M**. Methods for large-scale analysis of gene expression, protein localization and disruption phenotypes in *Saccharomyces cerevisiae*. *Methods Mol Cell Biol.* 1995;5: 298-308.
181. Heath CV, Copeland CS, Amberg DC, Del Priore V, **Snyder M**, Cole CN. Nuclear pore complex clustering and nuclear accumulation of poly(A)⁺ RNA associated with mutation of the *Saccharomyces cerevisiae* *RAT2/NUP120* gene. *J Cell Biol.* 1995;131: 1677-1697.
182. Sobel S, **Snyder M**. A highly divergent γ -tubulin is essential for cell growth and proper microtubule organization. *J Cell Biol.* 1995;131: 1775-1788.
183. Cid VJ, Duran A, del Rey F, **Snyder M**, Nombela C, Sanchez M. Molecular basis of cell integrity and morphogenesis in *Saccharomyces cerevisiae*. *Microbiol Rev.* 1995;59: 345-386.
184. Li O, Heath CV, Amberg DC, Dockendorff TC, Copeland CS, **Snyder M**, Cole CN. Mutation or deletion of the *Saccharomyces cerevisiae* *RAT3/NUP133* gene causes temperature-dependent nuclear accumulation of poly(A)⁺ RNA and constitutive clustering of nuclear pore complexes. *Mol Biol Cell.* 1995;6: 401-417.
185. Xie K, **Snyder M**. Two short autoepitopes on the nuclear dot antigen are similar to eptiopes encoded by Ebstein Barr Virus. *Proc Natl Acad Sci USA.* 1995;92: 1639-1643
186. **Snyder M**. The spindle pole body of yeast. *Chromosoma.* 1994;103: 369-380.
187. Costigan C, **Snyder M**. *SLK1*, a homolog of MAP kinase activators, mediates nutrient sensing independently of the yeast cAMP-dependent protein kinase pathway. *Mol Gen Genet.* 1994;243: 286-296.
188. Burns N, Grimwade B, Ross-Macdonald PB, Choi EY, Finberg K, Roeder GS, **Snyder M**. Large-scale analysis of gene expression, protein localization and gene disruption in *Saccharomyces cerevisiae*. *Genes Devel.* 1994;8: 1087-1105. (Featured in News and Views.)
189. Friedman H, **Snyder M**. Temperature-sensitive mutations in *PRG1*, a yeast proteasome-related gene cause defects in nuclear division and are suppressed by deletion of a mitotic cyclin gene. *Proc Natl Acad Sci USA.* 1994;91: 2031-2035.
190. Costigan C, Kolodrubetz D, **Snyder M**. *NHP6A* and *NHP6B*, which encode HMG1-like proteins, are candidates for downstream components of the yeast SLT2 mitogen-activated

protein kinase pathway. *Mol Cell Biol.* 1994;14: 2391-2403.

191. Page BD, Satterwhite LL, Rose MD, **Snyder M.** Localization of the Kar3 kinesin heavy chain-like protein requires the Cik1 interacting protein. *J Cell Biol.* 1994;124: 507-519.
192. Xie K, Lambie E, **Snyder M.** Nuclear dot antigens may specify transcriptional domains in the nucleus. *Mol Cell Biol.* 1993;13: 6170-6179.
193. Page BD, **Snyder M.** Chromosome segregation in yeast. *Ann Rev Microbiol.* 1993;47: 231-201.
194. Flescher EG, Madden K, **Snyder M.** Components required for cytokinesis are important for bud site selection in yeast. *J Cell Biol.* 1993;122: 373-386.
195. Granot D, **Snyder M.** Carbon source induces growth in stationary phase cells of yeast, independent of carbon source metabolism. *Yeast.* 1993;9: 465-479.
196. Copeland CS, **Snyder M.** Nuclear pore complex antigens delineate nuclear envelope dynamics in vegetative and conjugating *Saccharomyces cerevisiae*. *Yeast.* 1993;9: 235-249.
197. Yang CH, **Snyder M.** The nuclear mitotic apparatus protein (NuMA) is important in the establishment and maintenance of the bipolar mitotic spindle apparatus. *Mol Biol Cell.* 1992;3: 1259-1267.
198. Friedman H, Goebel M, **Snyder M.** A homolog of the MHC proteasome-related *RING10* gene is essential for yeast cell growth. *Gene.* 1992;122: 203-206.
199. Madden K, **Snyder M.** Specification of sites for polarized growth in *Saccharomyces cerevisiae* and the influence of external factors on site selection. *Mol Biol Cell.* 1992;3: 1025-1035.
200. Page BD, **Snyder M.** CIK1: a developmentally regulated spindle pole body-associated protein important for microtubule functions in *Saccharomyces cerevisiae*. *Genes Devel.* 1992;6: 1414-1429.
201. Mirzayan C, Copeland C, **Snyder M.** The *NUF1* gene encodes a coiled-coil related protein that is a potential component of the yeast nucleoskeleton. *J Cell Biol.* 1992;116: 1319-1332.
202. Yang C, Lambie EJ, **Snyder M.** NuMA: an unusually large coiled-coil protein in the mammalian nucleus. *J Cell Biol.* 1992;116: 1303-1317.
203. Costigan C, Gehrung S, **Snyder M.** A synthetic lethal screen identifies *SLK1*, a novel protein kinase homologue important in yeast cell morphogenesis and cell growth. *Mol Cell Biol.* 1992;12: 1162-1178.
204. Madden K, Costigan C, **Snyder M.** Cell polarity and morphogenesis in *Saccharomyces cerevisiae*. *Trends Cell Biol.* 1992;2: 22-29.
205. Granot D, **Snyder M.** Segregation of the nucleolus during mitosis in budding and fission yeast. *Cell Motility Cytoskel.* 1991;20: 47-54.
206. **Snyder M,** Gehrung S, Page BD. Studies concerning the temporal and genetic control of cell polarity in *Saccharomyces cerevisiae*. *J Cell Biol.* 1991;114: 515-532.

207. Padmanabha R, Gehrung S, **Snyder M**. The *KNS1* gene of *Saccharomyces cerevisiae* encodes a nonessential protein kinase homologue distantly related to members of the *CDC28/CDC2* gene family. *Mol Gen Genetics*. 1991;229: 1-9.
208. Granot D, **Snyder M**. Glucose induces cAMP-independent growth-related changes in stationary phase cells of *Saccharomyces cerevisiae*. *Proc Natl Acad Sci USA*. 1991;88: 5724-5728.
209. Gehrung S, **Snyder M**. The *SPA2* gene of *Saccharomyces cerevisiae* is important for pheromone-induced morphogenesis and efficient mating. *J Cell Biol*. 1990;111: 1451-1464.
210. Yang CH, Lambie EJ, Hardin J, Craft J, **Snyder M**. Higher order structure is present in the yeast nucleus: autoantibody probes demonstrate that the nucleolus lies opposite the spindle pole body. *Chromosoma*. 1989;98: 123-128.
211. **Snyder M**. The SPA2 protein of yeast localizes to sites of cell growth. *J Cell Biol*. 1989;108: 1419-1429.
212. Kolman CJ, **Snyder M**, Soll D. Genomic organization of tRNA and aminoacyl-tRNA synthetase genes for two amino acid families in *Saccharomyces cerevisiae*. *Genomics* 1988;3: 201-206.
213. **Snyder M**, Davis RW. *SP1*: a gene important for chromosome segregation and other mitotic functions in *S. cerevisiae*. *Cell*. 1988;54: 743-754.
214. **Snyder M**, Sapolsky R, Davis RW. Transcription interferes with elements important for chromosome maintenance. *Mol Cell Biol*. 1988;8: 2184-2194.
215. **Snyder M**, Elledge S, Sweetser D, Young RA, Davis RW. λ gt11: gene isolation with antibody probes and other applications. *Methods Enzymol*. 1987;154: 107-128.
216. **Snyder M**, Buchman AR, Davis RW. Bent DNA at a yeast autonomously replicating sequence. *Nature*. 1986;324: 87-89.
217. Davis RW, Vollrath D, Thomas M, Hieter P, **Snyder M**, Mann C, Fasullo M. Genome plasticity in *Saccharomyces cerevisiae*. 1986. UCLA Symposium series.
218. **Snyder M**, Elledge S, Davis RW. Rapid mapping of antigenic coding regions and creating insertion mutations in yeast genes by mini-Tn10 "transplason" mutagenesis. *Proc Natl Acad Sci USA*. 1986;83: 730-734.
219. Johnson LM, **Snyder M**, Chang LMS, Davis RW, Campbell JL. Isolation of the gene encoding yeast DNA polymerase I. *Cell*. 1985;43: 369-377.
220. Hieter P, Mann C, **Snyder M**, Davis RW. Mitotic stability of yeast chromosomes: a colony color assay that measures nondisjunction and chromosome loss. *Cell*. 1985;40: 381-392.
221. **Snyder M**, Davis RW. Screening λ gt11 expression libraries with antibody probes. *Hybridomas in the Biosciences and Medicine*, 1985. Timothy Springer, ed., Plenum Press, N.Y. p. 397-406.
222. **Snyder M**. Organization and expression of a cluster of *Drosophila* cuticle genes. Doctoral Dissertation, California Institute of Technology, Pasadena, CA, 1983.

223. **Snyder M**, Davidson N. Two gene families clustered in a small region of the *Drosophila* genome. *J Mol Biol.* 1983;166: 101-118.
224. **Snyder M**, Hunkapiller M, Yuen D, Silvert D, Fristrom J, Davidson N. Cuticle protein genes of *Drosophila*: structure, organization and evolution of four clustered genes. *Cell.* 1982;29: 1027-1040.
225. **Snyder M**, Kimbrell D, Hunkapiller M, Hill R, Fristrom J, Davidson N. A transposable element that splits the promoter region inactivates a *Drosophila* cuticle protein gene. *Proc Natl Acad Sci USA.* 1982;79: 7430-7434.
226. **Snyder M**, Hunkapiller M, Yuen D, Silvert D, Fristrom J, Davidson N. A cluster of *Drosophila* cuticle genes. In: ICN-UCLA Symposium on Developmental Biology Using Purified Genes XXIII, 1981. D Brown and F Fox, Eds., p.125-133.
227. **Snyder M**, Hirsh J, Davidson N. The cuticle genes of *Drosophila*: a developmentally regulated gene cluster. *Cell.* 1981;25: 165-177.
228. **Snyder M**, Drlica K. DNA Gyrase on the bacterial chromosome: DNA cleavage induced by oxolinic acid. *J Mol Biol.* 1979;131: 287-302.
229. Drlica K, **Snyder M**. Superhelical *Escherichia coli* DNA: relaxation by coumermycin. *J Mol Biol.* 1978;120: 145-154.